

Achievements and perspectives for the n_TOF facility at CERN

Alberto Mengoni
IAEA , Vienna


- The CERN n_TOF Facility
- Experimental campaigns in 2002-2004: status of the data analysis & results
- n_TOF-Phase 2

The n_TOF facility at CERN




somewhere around here

CERN accelerator Complex





Linac(s): up to 50 MeV PSB: up to 1 GeV PS: up to 24 GeV

The n_TOF facility at CERN



The real world

- n _TOF commissioned in 2001-2002




n_TOF beam characteristics

the neutron flux

2nd collimator $\phi=1.8$ cm
(capture mode)

Performance Report
CERN-INTC-2002-037, January 2003
CERN-SL-2002-053 ECT

$dN/d\ln E/7.e12$ protons



The neutron fluence in EAR-1

Energy range	Uncollimated [n/pulse/cm ²]	Capture mode [n/pulse]	Fission mode [n/pulse]
< 1 eV	2.0E+05	3.1E+05	2.0E+06
1 eV - 10 eV	2.7E+04	4.5E+04	2.9E+05
10 eV - 100 eV	2.9E+04	4.7E+04	3.1E+05
100 eV - 1000 eV	3.0E+04	5.1E+04	3.3E+05
1 eV - 1 keV	8.6E+04	1.4E+05	9.3E+05
1 keV - 10 keV	3.2E+04	5.4E+04	3.6E+05
10 keV - 100 keV	3.9E+04	7.1E+04	4.7E+05
100 keV - 1000 keV	1.1E+05	2.3E+05	1.5E+06
1 keV - 1 MeV	1.8E+05	3.5E+05	2.3E+06
1 MeV - 10 MeV	8.3E+04	2.4E+05	1.7E+06
10 MeV - 100 MeV	2.8E+04	7.2E+04	5.1E+05
> 100 MeV	4.4E+04	1.2E+05	5.6E+05
1 MeV - > 100 MeV	1.6E+05	4.4E+05	2.7E+06
Total	6.2E+05	1.2E+06	8.0E+06


Note: 1 pulse is 7E+12 protons. Collimated fluence (fission and capture modes) is integrated over the beam surface.

n_TOF basic parameters

proton beam momentum	20 GeV/c
intensity (dedicated mode)	7×10^{12} protons/pulse
repetition frequency	1 pulse/2.4s
pulse width	6 ns (rms)
n/p	300
lead target dimensions	80x80x60 cm ³
cooling & moderation material	H ₂ O
moderator thickness in the exit face	5 cm
neutron beam dimension in EAR-1 (capture mode)	2 cm (FWHM)

n_TOF beam characteristics


- Beam profile @ 187.5 m



MicroMegas detector

J Pancin, et al. (The n_TOF Collaboration)
NIMA 524 (2004) 102

J. Pancin et al. / Nuclear Instruments and Methods in Physics Research A 524 (2004) 102–114



FWHM \approx 2cm




Fig. 9. Horizontal (a), vertical (b), 30° (c) experimental and simulated projected profiles between 10 and 100 eV at 186 m.

n_TOF beam



energy resolution

Performance Report
CERN-INTC-2002-037, January 2003
CERN-SL-2002-053 ECT


Energy resolution @ 187.5 m (collimator for capture mode)

Neutron Energy	p-beam pulse width FWHM [cm]	moderation FWHM [cm]	$\Delta E/E$
1 eV	0.0	3.0	3.0E-04
10 eV	0.1	3.0	3.2E-04
100 eV	0.2	3.3	3.5E-04
1 keV	0.6	5.1	5.5E-04
10 keV	2.0	7.9	8.7E-04
30 keV	3.4	10.2	1.1E-03
100 keV	6.2	18.0	2.0E-03
1 MeV	19.5	34.1	4.2E-03
10 MeV	61.7	16.9	6.8E-03
100 MeV	195.0	14.5	2.1E-02

11.2 keV Fe 2.0x45mm C6D6



Monte Carlo RF: E =8-12keV




Chi2 / ndf = 0 / -9
t = -0.013 0
L = 0.02 0
A₂ = -0.163 0
A₃ = 14.45 0
A₄ = 0.163 0
A₅ = 13.5 0
t₀ = -0.07 0
A₁ = 450.3 0
A_n = -305 0

$$\frac{\Delta E}{E} = \frac{2}{L} \sqrt{\Delta L^2 + 1.91 \cdot E \cdot \Delta T^2}$$

(for example: 6×10^{-4} @ 1 keV)


n_TOF TAC for (n,γ) measurements

- 40 BaF₂ crystals
12 pentagons & 28 hexagons
15 cm crystal thickness
Carbon-fiber ¹⁰B-enriched capsules
- High detection efficiency: $\approx 100\%$
(to be compared with 5% of C₆D₆ or 0.1% of Moxon-Rae)
- Good energy resolution
(direct background suppression mechanisms
based on combined multiplicity and energy
deposition analysis)
- Full Monte Carlo simulations
all EM cascades
capture events for BG determination




TAC efficiency & MC simulations

The 4.9 eV resonances of ^{197}Au has been used as a reference sample and as a validation of the MC method for the estimation of the ε_{det} under different conditions on E_{sum} and m_{γ} .



TAC efficiency & MC simulations

The MC is capable of reproducing the E_{sum} spectrum for different multiplicities allowing to calculate very accurately the detection efficiency for the chosen analysis conditions




$$\sigma(\varepsilon_{\text{det}}) \leq 3\%$$

n_TOF TAC for (n,γ) measurements

- Structure mounted in April-04
- 4π geometry: end of May-04
- 1.5 month commissioning
- $\text{Au}(n,\gamma)$ & other standards



First measurement with a radioactive sample started in August 2004
 $^{237}\text{Np}(n,\gamma)$



n_TOF fission detectors

- 20x20 cm²
- Isobutane gas 7 mbar
- HV 500-600 V
- 3 mm between electrodes
- 1 anode (a few ns signal width)
- Electrode thickness: 1.5 μm (Mylar+Al)
- Deposit thickness : 100-300 μg/cm²
- Backing thickness : 0.1 μm (Al)
- : 1.5 μm (Mylar)

• Fission event identification: T2 in coincidence with T1



- Gas: Ar (90%) CF₄ (10%)
- Gas pressure : 720 mbar
- Electric field : 600 V/cm
- Gap pitch : 5 mm
- Electrode diameter : 12 cm
- Electrode thickness: 15 μm (Al)
- Deposit thickness : 125 μg/cm²
- Backing thickness : 100 μm (Al)
- Window thickness : 125 μm

Nuclear waste: TRU

(1000 MW_e LWR)

	Cm 238 2,4 h	Cm 239 3 h	Cm 240 27 d	Cm 241 32,8 d	Cm 242 162,94 d	Cm 243 29,1 a	Cm 244 18,10 a	Cm 245 8500 a	Cm 246 4730 a
Am 236 ? 3,7 m	Am 237 73,0 m	Am 238 1,63 h	Am 239 11,9 h	Am 240 50,8 h	Am 241 432,2 a	Am 242 141 a 16 h	Am 243 7370 a	Am 244 26 m 10,1 h	Am 245 2,05 h
Pu 235 25,3 m	Pu 236 2,858 a	Pu 237 45,2 d	Pu 238 87,74 a	Pu 239 2,411 · 10 ⁴ a	Pu 240 6563 a	Pu 241 14,35 a	Pu 242 3,750 · 10 ⁵ a	Pu 243 4,956 h	Pu 244 8,00 · 10 ⁷ a
Np 234 4,4 d	Np 235 396,1 d	Np 236 22,5 h 1,54 · 10 ⁵ a	Np 237 2,144 · 10 ⁶ a	Np 238 2,117 d	Np 239 2,355 d	Np 240 7,22 m 65 m	Np 241 13,9 m	Np 242 2,2 m 3,5 m	Np 243 1,85 m
U 233 1,592 · 10 ⁵ a	U 234 0,0055 a	U 235 0,7200 a	U 236 120 ns 2,342 · 10 ⁷ a	U 237 6,75 d	U 238 99,2745 a	U 239 23,5 m	U 240 14,1 h	U 242 16,8 m	
Pa 232 1,31 d	Pa 233 27,0 d	Pa 234 1,17 m 6,70 h	Pa 235 4,2 m	Pa 236 9,1 m	Pa 237 8,7 m	Pa 238 2,3 m	148	150	
Th 231 25,5 h	Th 232 100 a	Th 233 22,3 m	Th 234 24,10 d	Th 235 7,1 m	Th 236 37,5 m	Th 237 5,0 m			

²⁴⁴Cm
1.5 Kg/yr

²⁴¹Am: 11.6 Kg/yr
²⁴³Am: 4.8 Kg/yr

²³⁹Pu: 125 Kg/yr

²³⁷Np: 16 Kg/yr

LLFP
76.2 Kg/yr

LLFP

source: Actinide and Fission Product Partitioning and Transmutation – NEA (1999)


Nucleosynthesis: the s-process

- 1/2 of the elements above Fe are produced by the s-process
- The astrophysical sites of the s-process are:
 - He burning in intermediate/massive stars
 - Low-mass AGB's
- There exists a direct correlation between the neutron capture cross section and the abundance ($\sigma(n, \gamma) \cdot N = const.$)
- The neutron capture cross sections are key ingredients for s-process nucleosynthesis

The canonical s-process



Cu			62Cu 9.74 m	63Cu 69.17	64Cu 12.7 h	
Ni		60Ni 26.223	61Ni 1.140	62Ni 3.634	63Ni 100 a	
Co		58Co 70.86 d	59Co 100	60Co 5.272 a	61Co 1.65 h	
Fe	56Fe 91.72	57Fe 2.2	58Fe 0.28	59Fe 44.503 d	60Fe 1.5 10 ⁶ a	61Fe 6 m

Solar system elemental abundances



Nucleosynthesis: the s-process & the r-process residuals

$$N_r = N_{\text{solar}} - N_s$$



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments 2002-4

- **M**easurements of neutron cross sections relevant for Nuclear Waste Transmutation and related Nuclear Technologies
 - Th/U fuel cycle (capture & fission)
 - Transmutation of MA (capture & fission)
 - Transmutation of FP (capture)
- **C**ross sections relevant for Nuclear Astrophysics
 - s-process: branchings
 - s-process: presolar grains
- **N**eutrons as probes for fundamental Nuclear Physics
 - Nuclear level density & n-nucleus interaction

n_TOF experiments 2002-4

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$ ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$ ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

- data analysis completed, results published
- data analysis completed, paper in preparation
- data analysis in progress

The n_TOF-Ph2 experiments

Capture measurements

Mo, Ru, Pd stable isotopes

r-process residuals calculation
isotopic patterns in SiC grains

Fe, Ni, Zn, and Se (stable isotopes)
 ^{79}Se

s-process nucleosynthesis in massive stars
accurate nuclear data needs for structural materials

$A \approx 150$ (isotopes varii)

s-process branching points
long-lived fission products

$^{234,236}\text{U}$, $^{231,233}\text{Pa}$

Th/U nuclear fuel cycle

$^{235,238}\text{U}$

standards, conventional U/Pu fuel cycle

$^{239,240,242}\text{Pu}$, $^{241,243}\text{Am}$, ^{245}Cm

incineration of minor actinides

(*) approved by CERN Scientific Committee (planned for execution in 2007)

The n_TOF-Ph2 experiments

Fission measurements

MA

ADS, high-burnup, GEN-IV reactors

$^{235}\text{U}(n,f)$ with $p(n,p')$

new $^{235}\text{U}(n,f)$ cross section standard

$^{234}\text{U}(n,f)$

study of vibrational resonances at the fission barrier

Other measurements

$^{147}\text{Sm}(n,\alpha)$, $^{67}\text{Zn}(n,\alpha)$, $^{99}\text{Ru}(n,\alpha)$

p-process studies

$^{58}\text{Ni}(n,p)$, other $(n,lc p)$

gas production in structural materials

Al, V, Cr, Zr, Th, $^{238}\text{U}(n,lc p)$

structural and fuel material for ADS
and other advanced nuclear reactors

He, Ne, Ar, Xe


low-energy nuclear recoils
(development of gas detectors)

$n+\text{D}_2$


neutron-neutron scattering length

NEW target design


xz-squared target (40x40x55) with
5cm-thick cylinder moderator
containers




H₂O





D₂O




D₂O cooling
(1cm Pb + 1cm H₂O)



NEW: target design proposal



The second n_TOF beam line & EAR-2



Flight-path length : ~20 m
at 90° respect to p-beam direction
expected neutron flux enhancement: ~ 100
drastic reduction of the t_0 flash

EAR-2: Optimized sensitivity

Improvements (ex: ^{151}Sm case)	consequences for sample mass
■ sample mass / 3 s/bkgd=1	✓ 50 mg
■ use BaF_2 TAC $\epsilon \times 10$	✓ 5 mg
■ use D_2O $\Phi_{30} \times 5$	■ 1 mg
■ use 20 m flight path $\Phi_{30} \times 100$	■ 10 μg

boosts sensitivity by a factor of 5000 !



→ problems of sample production and safety issues relaxed

The n_TOF Collaboration

U.Abbondanno¹⁴, G.Aerts⁷, H.Álvarez²⁴, F.Alvarez-Velarde²⁰, S.Andriamonje⁷, J.Andrzejewski³³, P.Assimakopoulos⁹, L.Audouin⁵, G.Badurek¹, P.Baumann⁶, F. Bečvář³¹, J.Benlliure²⁴, E.Berthoumieux⁷, F.Calviño²⁵, D.Cano-Ott²⁰, R.Capote²³, A.Carrillo de Albornoz³⁰, P.Cennini⁴, V.Chepel⁷, E.Chiaveri⁴, N.Colonna³, G.Cortes²⁵, D.Cortina²⁴, A.Couture²⁹, J.Cox²⁹, S.David⁵, R.Dolfini¹⁵, C.Domingo-Pardo²¹, W.Dridi⁷, I.Duran²⁴, M.Embid-Segura²⁰, L.Ferrant⁵, A.Ferrari⁴, R.Ferreira-Marques¹⁷, L.Fitzpatrick⁴, H.Frais-Koelbl³, K.Fujii¹³, W.Furman¹⁸, C.Guerrero²⁰, I.Goncalves³⁰, R.Gallino³⁶, E.Gonzalez-Romero²⁰, A.Goverdovski¹⁹, F.Gramegna¹², E.Griesmayer³, F.Gunsing⁷, B.Haas³², R.Haight²⁷, M.Heil⁸, A.Herrera-Martinez⁴, M.Igashira³⁷, S.Isaev⁵, E.Jericha¹, Y.Kadi⁴, F.Käppeler⁸, D.Karamanis⁹, D.Karadimos⁹, M.Kerveno⁶, V.Ketlerov¹⁹, P.Koehler²⁸, V.Konovalov¹⁸, E.Kossionides³⁹, M.Krtička³¹, C.Lamboudis¹⁰, H.Leeb¹, A.Lindote¹⁷, I.Lopes¹⁷, M.Lozano²³, S.Lukic⁶, J.Marganec³³, L.Marques³⁰, S.Marrone¹³, P.Mastinu¹², A.Mengoni⁴, P.M.Milazzo¹⁴, C.Moreau¹⁴, M.Mosconi⁸, F.Neves¹⁷, H.Oberhummer¹, S.O'Brien²⁹, M.Oshima³⁸, J.Pancin⁷, C.Papachristodoulou⁹, C.Papadopoulos⁴⁰, C.Paradela²⁴, N.Patronis⁹, A.Pavlik², P.Pavlopoulos³⁴, L.Perrot⁷, R.Plag⁸, A.Plompen¹⁶, A.Plukis⁷, A.Poch²⁵, C.Pretel²⁵, J.Quesada²³, T.Rauscher²⁶, R.Reifarth²⁷, M.Rosetti¹, C.Rubbia⁵, G.Rudolf⁶, P.Rullhusen¹⁶, J.Salgado³⁰, L.Sarchiapone⁴, C.Stephan⁵, G.Tagliente¹³, J.L.Tain²¹, L.Tassan-Got⁵, L.Tavora³⁰, R.Terlizzi¹³, G.Vannini³⁵, P.Vaz³⁰, A.Ventura¹¹, D.Villamarin²⁰, M.C.Vincente²⁰, V.Vlachoudis⁴, R.Vlastou⁴⁰, F.Voss⁸, H.Wendler⁴, M.Wiescher²⁹, K.Wisshak⁸

40 Research Institutions
120 researchers

The End

PS: all quoted documents are available online at

www.cern.ch/ntof

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

U Abbondanno et al. (The n_TOF Collaboration)
Phys. Rev. Lett. **93** (2004), 161103



MACS-30 = 3100 ± 160 mb

Capture

¹⁵¹Sm

^{204,206,207,208}Pb, ²⁰⁹Bi

²³²Th

^{24,25,26}Mg

^{90,91,92,94,96}Zr, ⁹³Zr

¹³⁹La

^{186,187,188}Os

^{233,234}U

²³⁷Np, ²⁴⁰Pu, ²⁴³Am

Fission

^{233,234,235,236,238}U

²³²Th

²⁰⁹Bi

²³⁷Np


^{241,243}Am, ²⁴⁵Cm



n_TOF experiments

U Abbondanno et al. (The n_TOF Collaboration)
Phys. Rev. Lett. **93** (2004), 161103 &

S Marrone et al. (The n_TOF Collaboration)
Phys. Rev. C **73** 03604 (2006)



Capture

¹⁵¹Sm

^{204,206,207,208}Pb, ²⁰⁹Bi

²³²Th

^{24,25,26}Mg

^{90,91,92,94,96}Zr, ⁹³Zr

¹³⁹La

^{186,187,188}Os

^{233,234}U

²³⁷Np, ²⁴⁰Pu, ²⁴³Am

Fission

^{233,234,235,236,238}U

²³²Th

²⁰⁹Bi

²³⁷Np


^{241,243}Am, ²⁴⁵Cm



n_TOF experiments

U Abbondanno et al. (The n_TOF Collaboration)
Phys. Rev. Lett. **93** (2004), 161103 &

S Marrone et al. (The n_TOF Collaboration)
Phys. Rev. C **73** 03604 (2006)



$$\begin{aligned} \langle D_0 \rangle &= 1.49 \pm 0.07 \text{ eV} \\ S_0 &= (3.87 \pm 0.33) \times 10^{-4} \\ R_1 &= 3575 \pm 210 \text{ b} \end{aligned}$$

Capture

¹⁵¹Sm

^{204,206,207,208}Pb, ²⁰⁹Bi

²³²Th

^{24,25,26}Mg

^{90,91,92,94,96}Zr, ⁹³Zr

¹³⁹La

^{186,187,188}Os

^{233,234}U

²³⁷Np, ²⁴⁰Pu, ²⁴³Am

Fission

^{233,234,235,236,238}U

²³²Th

²⁰⁹Bi

²³⁷Np

^{241,243}Am, ²⁴⁵Cm



n_TOF experiments

U Abbondanno et al. (The n_TOF Collaboration)
Phys. Rev. Lett. **93** (2004), 161103
&

S Marrone et al. (The n_TOF Collaboration)
Phys. Rev. C **73** 03604 (2006)

for nuclear data
evaluators:
all infos available in
refereed journal
publications
&
on the n_TOF website
www.cern.ch/ntof

TABLE IX. The ¹⁵¹Sm(*n,γ*) cross section in the unresolved resonance region from 1 keV to 1 MeV.

Energy bin (keV)	$\sigma_{(n,\gamma)}$ (b)	Uncertainty (%)		
		Stat.	Syst.	Tot.
1–1.2	24.52	0.8	4.4	4.5
1.2–1.5	23.68	0.8	4.3	4.4
1.5–1.75	21.94	1.0	4.2	4.3
1.75–2	19.76	1.2	4.2	4.3
2–2.5	15.43	1.1	4.1	4.3
2.5–3	15.36	1.3	4.1	4.3
3–4	12.78	1.2	4.1	4.3
4–5	10.04	1.4	4.1	4.3
5–7.5	8.91	2.1	2.9	3.6
7.5–10	5.85	3.0	3.1	4.3
10–12.5	5.38	3.9	2.9	4.8
12.5–15	4.26	4.9	3.2	5.8
15–20	3.82	3.8	3.2	4.9
20–25	3.52	4.6	3.5	5.8
25–30	3.13	4.5	3.1	5.5
30–40	2.69	4.4	3.2	5.5
40–50	2.17	4.8	3.4	5.9
50–60	1.90	5.2	3.3	6.2
60–80	1.66	4.1	3.6	5.5
80–100	1.30	5.1	4.6	6.9

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}, ^{209}\text{Bi}$

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}, ^{93}\text{Zr}$

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

$^{237}\text{Np}, ^{240}\text{Pu}, ^{243}\text{Am}$

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi



^{237}Np

$^{241,243}\text{Am}, ^{245}\text{Cm}$

n_TOF experiments

C Domingo-Pardo, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004
&
accepted for publication in PRC (in press)

$^{207}\text{Pb}(n,\gamma)$



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}, ^{209}\text{Bi}$

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}, ^{93}\text{Zr}$

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

$^{237}\text{Np}, ^{240}\text{Pu}, ^{243}\text{Am}$

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}, ^{245}\text{Cm}$

n_TOF experiments

C Domingo-Pardo, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004 &
accepted for publication in PRC (in press)

$^{207}\text{Pb}(n,\gamma)$



substantial disagreement for $E_n > 45$ keV

Capture

¹⁵¹Sm

204,206,207,208Pb, ²⁰⁹Bi

²³²Th

24,25,26Mg

90,91,92,94,96Zr, ⁹³Zr

¹³⁹La

186,187,188Os

^{233,234}U

²³⁷Np, ²⁴⁰Pu, ²⁴³Am

Fission

^{233,234,235,236,238}U

²³²Th

²⁰⁹Bi

²³⁷Np

^{241,243}Am, ²⁴⁵Cm

n_TOF experiments

C Domingo-Pardo, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004
&
accepted for publication in PRC (in press)

²⁰⁷Pb(n,γ)

TABLE II: Resonance parameters and radiative kernels from the analysis of the ²⁰⁷Pb(n,γ) data measured at n_TOF^a.

E_0 (eV)	l	J	Γ_n (meV)	Γ_γ (meV)	$g\Gamma_\gamma\Gamma_n/\Gamma$ (meV)
3064.700(3)	1	2	111.0(8)	145.0(9)	78.6(9)
10190.80(4)	1	2	656(50)	145.2(12)	149(14)
16172.80(10)	1	2	1395(126)	275(3)	287(30)
29396.1	1	2	16000	189(7)	234(9)
30485.9(5)	1	1	608(45)	592(50)	225(30)
37751(3)	1	1	50×10^3	843(40)	620(30)
41149(46)	0	1	1.220×10^6	3970(160)	2970(120)
48410(2)	1	2	1000	230(20)	235(20)
82990(12)	1	2	29×10^3	360(30)	444(30)
90228(24)	1	1	272×10^3	1615(100)	1200(80)
127900	1	1	613×10^3	1939(150)	1449(120)
130230	1	1	87×10^3	900(80)	675(60)
181510(6)	0	1	57.3×10^3	14709(500)	8780(300)
254440	2	3	111×10^3	1219(90)	2110(150)
256430	0	1	1.66×10^6	12740(380)	9482(280)
317000	0	1	850×10^3	10967(480)	8120(350)

^aOrbital angular momenta l and resonance spins J are from Ref. [17].

3% accuracy
of the capture kernel

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}, ^{209}\text{Bi}$

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}, ^{93}\text{Zr}$

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

$^{237}\text{Np}, ^{240}\text{Pu}, ^{243}\text{Am}$

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}, ^{245}\text{Cm}$

n_TOF experiments

C Domingo-Pardo, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004
&
submitted for publication to PRC, October 2006

$^{204}\text{Pb}(n,\gamma)$



Very low neutron sensitivity of capture γ -ray
detection systems & high resolution

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}, ^{209}\text{Bi}$

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}, ^{93}\text{Zr}$

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

$^{237}\text{Np}, ^{240}\text{Pu}, ^{243}\text{Am}$

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}, ^{245}\text{Cm}$

n_TOF experiments

C Domingo-Pardo, et al. - The n_TOF Collaboration
 ND2004 Conference, Santa Fe, NM – Sept. 2004
 &
 submitted for publication to PRC, October 2006

$^{204}\text{Pb}(n,\gamma)$

TABLE IV: Average neutron capture cross section for ^{204}Pb .

E_o (eV)	l	J	Γ_γ (meV)	$\Delta\Gamma_\gamma$ (%)	Γ_n (meV)	K_r (meV)	ΔK_r (%)
480.3	1	1/2	1.33	4	3.0	0.92 ^a	2.7
1333.8	1	1/2	105	4	46.3 ^b	32.1 ^a	1.3
1687.1	0	1/2	1029	0.7	3340	787 ^a	0.5
2481.0	0	1/2	514	1.1	5470	470 ^a	1.0
2600.0						8.35	6
2707.1	1	3/2	31.2	9	11.5	16.8	2
3187.9	0	1/2	316	10	1.7	1.69	0.1
3804.9	1	1/2	280	8	66.4	53.7	1.6
4284.1	1	3/2	111	9	24.0	39.4	1.7
4647.5						2.57	9
4719.4	1	3/2	41.2	5	95.0	57.5	3
5473.2	1	1/2				79.0	1.6
5561.4		(1/2)	1.03	10	1.9	0.67	6.4
6700.5	0	1/2	312	3	4540	292	3
7491.0						19.0	0.5
8357.4	0	1/2	1286	1.9	45000	1250	1.9
8422.9						11.3	7
8949.6						22.9	3
9101.0		(1/2)	193	8	150	84.4	4
9649.3	0	1/2	1076	2	7860	946	2
10254						37.0	8
11366	1	3/2	39.0	10	226	66.5	9
11722						22.8	9
12147						54.4	8

E_{low} (keV)	E_{high} (keV)	Cross section (barn)	Statistical uncertainty ^a (%)
88.210	92.404	0.059	9
92.404	96.748	0.059	5
96.748	101.406	0.058	11
101.406	106.408	0.057	8
106.408	111.790	0.057	7
111.790	117.591	0.056	8
117.591	123.855	0.056	7
123.855	130.634	0.055	7
130.634	137.985	0.054	6
137.985	145.974	0.054	6
145.974	154.678	0.053	6
154.678	164.185	0.053	7
164.185	174.596	0.052	7
174.596	186.030	0.051	6
186.030	198.625	0.051	5
198.625	212.544	0.050	5
212.544	227.981	0.049	5
227.981	245.162	0.049	5
245.162	264.363	0.048	4
264.363	285.911	0.047	4
285.911	310.207	0.046	4
310.207	337.739	0.046	4
337.739	369.107	0.045	4
369.107	405.060	0.044	4
405.060	443.512	0.043	3

^aThis value has to be added in quadrature with the overall systematic uncertainty of 10%.

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi



^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

C Domingo-Pardo, et al. (The n_TOF Collaboration)
Phys. Rev. C 74, 025807 (2006)

$^{209}\text{Bi}(n,\gamma)$



Very low neutron sensitivity of capture γ -ray detection systems & high resolution

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi



^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

$^{209}\text{Bi}(n,\gamma)$

C Domingo-Pardo, et al. (The n_TOF Collaboration)
Phys. Rev. C 74, 025807 (2006)



Very low neutron sensitivity of capture γ -ray detection systems & high resolution

Capture

¹⁵¹Sm

204,206,207,208Pb, ²⁰⁹Bi

²³²Th

24,25,26Mg

90,91,92,94,96Zr, ⁹³Zr

¹³⁹La

186,187,188Os

^{233,234}U

²³⁷Np, ²⁴⁰Pu, ²⁴³Am

Fission

^{233,234,235,236,238}U

²³²Th

²⁰⁹Bi

²³⁷Np

^{241,243}Am, ²⁴⁵Cm

n_TOF experiments

C Domingo-Pardo, et al. (The n_TOF Collaboration)
Phys. Rev. C **74**, 025807 (2006)

²⁰⁹Bi(n,γ)

NEW MEASUREMENT OF NEUTRON CAPTURE . . .

PHYSICAL REVIEW C **74**, 025807 (2006)

TABLE II. Resonance parameters^a and radiative kernels^b for ²⁰⁹Bi.

E_o (eV)	l	J	Γ_n (meV)	Γ_γ (meV)	$g\Gamma_\gamma\Gamma_n / \Gamma$ (meV)
801.6(1)	0	5	4309(145)	33.3(12)	18.2(6)
2323.8(6)	0	4	17888(333)	26.8(17)	12.0(8)
3350.83(4)	1	5	87(9)	18.2(3)	9.5(2)
4458.74(2)	1	5	173(13)	23.2(22)	11.3(11)
5114.0(3)	0	5	5640(270)	65(2)	35.3(11)
6288.59(2)	1	4	116(18)	17.0(17)	6.7(7)
6525.0(3)	1	3	957(100)	25.3(14)	8.6(5)
9016.8(4)	1	6	408(77)	21.1(14)	13.0(9)
9159.20(7)	1	5	259(45)	21.4(21)	10.9(11)
9718.910(1)	1	4	104(22)	74(7)	19.5(21)
9767.2(3)	1	3	900(114)	90(8)	28.7(26)
12098					65(4) ^c
15649.8(1.0)	1	5	1000	47(4)	20.2(17)
17440.0(1.3)	1	6	1538(300)	32(3)	20.4(18)
17839.5(9)	1	5	464(181)	43(4)	21.7(20)
20870	1	5	954(227)	34.4(33)	18.3(17)
21050	1	4	7444(778)	33(3)	14.8(13)
22286.0(9)	1	5	181(91)	33.6(32)	15.1(15)
23149.1(1.3)	1	6	208(154)	25.3(25)	14.7(15)

^aAngular orbital momenta, l , resonance spins J , and neutron widths, Γ_n , are mainly from Refs. [27,28].

^bUncertainties are given as 18.2(6)≡18.2±0.6.

^cThis area corresponds to the sum of the areas of the broad s -wave resonance at the indicated energy, plus two p -wave resonances at 12.092 and 12.285 keV.

16% higher MACS for kT = 5-8 keV
81% r-process abundance for ²⁰⁹Bi

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Günsing, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004



Low PS duty-cycle favours measurements
on radioactive samples

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm




$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Gunsing, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004

&
G Aerts et al. (The n_TOF Collaboration)
Phys. Rev. C 73, 054610 (2006)



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Gunsing, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004

&
G Aerts et al. (The n_TOF Collaboration)
Phys. Rev. C 73, 054610 (2006)

TABLE II. Different components of estimated systematic or correlated uncertainty in the measured cross section.

Component	Uncertainty (%)
PHWT	0.5
Normalization	0.5
Background	2.5
Flux shape	2.0
Total	3.3

For $E_n = 4 \text{ keV}$ up to 1 MeV full dataset is available on the PRC publication

E_{low} (keV)	E_{high} (keV)	Cross section (b)	Uncertainty (b)
3.994	4.482	0.958	0.020
4.482	5.028	1.281	0.021
5.028	5.642	1.097	0.016
5.642	6.331	1.004	0.014
6.331	7.103	0.912	0.013
7.103	7.970	0.919	0.013
7.970	8.942	0.848	0.013
8.942	10.033	0.817	0.012
10.033	11.257	0.800	0.012
11.257	12.631	0.787	0.012
12.631	14.172	0.761	0.012
14.172	15.902	0.729	0.011
15.902	17.842	0.685	0.011
17.842	20.019	0.613	0.010
20.019	22.461	0.641	0.010
22.461	25.202	0.566	0.009
25.202	28.277	0.545	0.009
28.277	31.728	0.513	0.008
31.728	35.599	0.497	0.009
35.599	39.943	0.468	0.009
39.943	44.816	0.456	0.008
44.816	50.285	0.413	0.007
50.285	56.421	0.365	0.006
56.421	63.305	0.346	0.006
63.305	71.029	0.318	0.006
71.029	79.696	0.275	0.005
79.696	89.421	0.248	0.005
89.421	100.332	0.229	0.005
100.332	112.574	0.220	0.004
112.574	126.310	0.204	0.004
126.310	141.722	0.192	0.004

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Gensing, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004



RRR region analysis in progress

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Gunsing, et al. - The n_TOF Collaboration analysis in progress



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,\gamma)$

n_TOF experiments

F Gunsing, et al. - The n_TOF Collaboration analysis in progress



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$


^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments



Very low neutron sensitivity of capture γ -ray
detection systems & high resolution

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$



^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments



Source: P Koehler & S O'Brien

Capture & transmission data (from ORELA)
analyzed simultaneously

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np




$^{241,243}\text{Am}$, ^{245}Cm

$^{96}\text{Zr}(n,\gamma)$

20% reduction
in the capture
strength
(average)

n_TOF experiments

C Moreau, et al.
ND2004 Conference, Santa
G Tagliente et al.



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

$^{90}\text{Zr}(n,\gamma)$

n_TOF experiments

C Moreau, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – September 2004
G Tagliente et al. (The n_TOF Collaboration)
NIC-IX, CERN, June 2006



Capture

- ^{151}Sm
- $^{204,206,207,208}\text{Pb}$, ^{209}Bi
- ^{232}Th
- $^{24,25,26}\text{Mg}$
- $^{90,91,92,94,96}\text{Zr}$ **^{93}Zr**
- ^{139}La
- $^{186,187,188}\text{Os}$
- $^{233,234}\text{U}$
- ^{237}Np , ^{240}Pu , ^{243}Am



Fission

- $^{233,234,235,236,238}\text{U}$
- ^{232}Th
- ^{209}Bi
- ^{237}Np
- $^{241,243}\text{Am}$, ^{245}Cm



$^{93}\text{Zr}(n,\gamma)$: raw data

n_TOF experiments



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

$^{139}\text{La}(n,\gamma)$

R Terlizzi, et al. (The n_TOF Collaboration)


CGS12

Notre Dame, IN, USA

AIP Conference Proceedings 819

&

submitted for publication to PRC, October 2006



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

$^{139}\text{La}(n,\gamma)$

R Terlizzi, et al. (The n_TOF Collaboration)


CGS12

Notre Dame, IN, USA

AIP Conference Proceedings 819

&

submitted for publication to PRC, October 2006



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th


^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

$^{139}\text{La}(n,\gamma)$



Remarkable energy resolution and background conditions have allowed to determine the resonance parameters up to 9 keV

$$R_I = 10.8 \pm 1.0 \text{ barn}$$

average γ -widths:

$$s\text{-waves} = 50.7 \pm 5.4 \text{ meV}$$

$$p\text{-waves} = 33.6 \pm 6.9 \text{ meV}$$

$$\langle D_0 \rangle = 252 \pm 22 \text{ eV}$$

$$S_0 = (0.82 \pm 0.05) \times 10^{-4} \quad S_1 = (0.55 \pm 0.04) \times 10^{-4}$$

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$


^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments



Very low neutron sensitivity of capture γ -ray detection systems & high resolution

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

M Mosconi, *et al.* – (The n_TOF Collaboration)
NIC-IX, CERN, Geneva – June 2006
analysis completed - paper in preparation

MACS-30

BrB81	438 ± 30 mb
WiM82	418 ± 16 mb
n_TOF	384 ± 17 mb



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm

n_TOF experiments

M Mosconi, *et al.* – (The n_TOF Collaboration)
NIC-IX, CERN, Geneva – June 2006
analysis completed - paper in preparation

MACS-30

BrB81	919 ± 28 mb
WiM82	874 ± 28 mb
n_TOF	940 ± 18 mb



Capture

^{151}Sm
 $^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

$^{233}\text{U}(n,\gamma)$

W Dridi, E Berthoumieux, *et al.*, (Dec. 2004)



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{233}\text{U}(n,\gamma)$

n_TOF experiments

W Dridi, E Berthoumieux, *et al.*, CEA/Saclay
Paper in preparation (October 2006)



n_TOF TAC in operation: capture & fission discrimination

The n_TOF Collaboration

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm




n_TOF experiments

W Dridi, E Berthoumieux, et al. (The n_TOF Collaboration)
PHYSOR-2006, Vancouver, September 2006
full paper in preparation

$^{234}\text{U}(n,\gamma)$

Figure 3: Neutron capture on ^{234}U yield in the thermal region and for the first resonance obtained in the present experiment.



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

W Dridi, E Berthoumieux, et al. (The n_TOF Collaboration)
PHYSOR-2006, Vancouver, September 2006
full paper in preparation

$^{234}\text{U}(n,\gamma)$



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

W Dridi, E Berthoumieux, et al. (The n_TOF Collaboration)
PHYSOR-2006, Vancouver, September 2006
full paper in preparation

$^{234}\text{U}(n,\gamma)$



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

W Dridi, E Berthoumieux, et al. (The n_TOF Collaboration)
PHYSOR-2006, Vancouver, September 2006
full paper in preparation

$^{234}\text{U}(n,\gamma)$



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

W Dridi, E Berthoumieux, et al. (The n_TOF Collaboration)
PHYSOR-2006, Vancouver, September 2006
full paper in preparation

$^{234}\text{U}(n,\gamma)$



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

D Cano-Ott, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

C Guerero, D Cano-Ott, et al. - The n_TOF Collaboration
PHYSOR 2006, Vancouver, September 2006

n_TOF ^{237}Np $\sigma(n,\gamma)$ compared to Evaluated Data Libraries



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np



$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

C Guerero, D Cano-Ott, et al. - The n_TOF Collaboration
PHYSOR 2006, Vancouver, September 2006

^{237}Np experimental Yield fitted with SAMMY



Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

C Guerero, D Cano-Ott, et al. - The n_TOF Collaboration
PHYSOR 2006, Vancouver, September 2006



$\text{RK}_{\text{n_TOF}}$ on average 3% below the RK_{JENDL} and 6% below the RK_{ENDF}

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

D Cano-Ott, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

C Guerero, D Cano-Ott, et al. - The n_TOF Collaboration
PHYSOR 2006, Vancouver, September 2006

n_TOF ^{240}Pu $\sigma(n,\gamma)$ compared to Evaluated Data Libraries



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

C Guerero, D Cano-Ott, et al. - The n_TOF Collaboration
PHYSOR 2006, Vancouver, September 2006



$\text{RK}_{\text{n_TOF}}$ is on average 9% smaller than RK_{JENDL} and 7% smaller than RK_{ENDF} .

Capture

- ^{151}Sm
- $^{204,206,207,208}\text{Pb}$, ^{209}Bi
- ^{232}Th
- $^{24,25,26}\text{Mg}$
- $^{90,91,92,94,96}\text{Zr}$, ^{93}Zr
- ^{139}La
- $^{186,187,188}\text{Os}$
- $^{233,234}\text{U}$
- ^{237}Np , ^{240}Pu , ^{243}Am


Fission

- $^{233,234,235,236,238}\text{U}$
- ^{232}Th
- ^{209}Bi
- ^{237}Np
- $^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

D Cano-Ott, et al. - The n_TOF Collaboration
ND2004 Conference, Santa Fe, NM – Sept. 2004



n_TOF TAC in operation

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{234}\text{U}(n,f)$

n_TOF experiments

PPACs & FIC-0 (2003)



An unprecedented wide energy range can be explored at n_TOF in a single experiment

Capture

- ^{151}Sm
- $^{204,206,207,208}\text{Pb}$, ^{209}Bi
- ^{232}Th
- $^{24,25,26}\text{Mg}$
- $^{90,91,92,94,96}\text{Zr}$, ^{93}Zr
- ^{139}La
- $^{186,187,188}\text{Os}$
- $^{233,234}\text{U}$
- ^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$


- ^{232}Th
- ^{209}Bi
- ^{237}Np
- $^{241,243}\text{Am}$, ^{245}Cm



$^{234}\text{U}(n,f)$

n_TOF experiments

PPACs & FIC-0 (2003)



High-resolution data up to high(er) energies

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{234}\text{U}(n,f)$

n_TOF experiments

PPACs & FIC-0 (2003)



High-resolution data up to high(er) energies

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

FIC-0 (2003)



An unprecedented wide energy range can be explored at n_TOF in a single experiment

Capture

- ^{151}Sm
- $^{204,206,207,208}\text{Pb}$, ^{209}Bi
- ^{232}Th
- $^{24,25,26}\text{Mg}$
- $^{90,91,92,94,96}\text{Zr}$, ^{93}Zr
- ^{139}La
- $^{186,187,188}\text{Os}$
- $^{233,234}\text{U}$
- ^{237}Np , ^{240}Pu , ^{243}Am

Fission


$^{233,234,235,236,238}\text{U}$

- ^{232}Th
- ^{209}Bi
- ^{237}Np
- $^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

FIC-1 (2003)



An unprecedented wide energy range can be explored at n_TOF in a single experiment

Capture

^{151}Sm
 $^{204,206,207,208}\text{Pb}$, ^{209}Bi
 ^{232}Th
 $^{24,25,26}\text{Mg}$
 $^{90,91,92,94,96}\text{Zr}$, ^{93}Zr
 ^{139}La
 $^{186,187,188}\text{Os}$
 $^{233,234}\text{U}$
 ^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{232}\text{Th}(n,f)$

n_TOF experiments

PPAC detectors



An unprecedented wide energy range can be explored at n_TOF in a single experiment

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{237}\text{Np}(n,f)$

n_TOF experiments

FIC-0 (2003)



Higher fission σ_f in the sub-threshold region

Capture

- ^{151}Sm
- $^{204,206,207,208}\text{Pb}$, ^{209}Bi
- ^{232}Th
- $^{24,25,26}\text{Mg}$
- $^{90,91,92,94,96}\text{Zr}$, ^{93}Zr
- ^{139}La
- $^{186,187,188}\text{Os}$
- $^{233,234}\text{U}$
- ^{237}Np , ^{240}Pu , ^{243}Am

Fission


- $^{233,234,235,236,238}\text{U}$
- ^{232}Th
- ^{209}Bi
- ^{237}Np
- $^{241,243}\text{Am}$, ^{245}Cm



$^{237}\text{Np}(n,f)$

n_TOF experiments

PPACs (2003)



Higher fission x-section in the sub-threshold region

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi


^{237}Np

$^{241,243}\text{Am}$, ^{245}Cm



n_TOF experiments

FIC-1 (2003)



High-resolution data up to high(er) energies

Capture

^{151}Sm

$^{204,206,207,208}\text{Pb}$, ^{209}Bi

^{232}Th

$^{24,25,26}\text{Mg}$

$^{90,91,92,94,96}\text{Zr}$, ^{93}Zr

^{139}La

$^{186,187,188}\text{Os}$

$^{233,234}\text{U}$

^{237}Np , ^{240}Pu , ^{243}Am

Fission

$^{233,234,235,236,238}\text{U}$

^{232}Th

^{209}Bi

^{237}Np


$^{241,243}\text{Am}$, ^{245}Cm



$^{238}\text{U}(n,f)/^{238}\text{U}(n,f)$

n_TOF experiments

FIC-0 (2003)



15% lower U8/U5 ratio at high energies


back

Capture studies: Mo, Ru and Pd

Motivations:

- Accurate determination of the r-process abundances (r-process residuals) from observations
- SiC grains carry direct information on s-process efficiencies in individual AGB stars. Abundance ratios in SiC grains strongly depend on available capture cross sections data.

$$N_r = N_{\text{solar}} - N_s$$




Capture studies: Mo, Ru and Pd

- Setup: The **n_TOF TAC** in EAR-1 (a few cases with C₆D₆ if larger neutron scattering)
- All samples are stable and non-hazardous
- Metal samples preferable (oxides acceptable)

Estimated # of protons
 $20 \times 5 \times 10^{16} = 10^{18}$


Cd 97 3 s	Cd 98 9.2 s	Cd 99 16 s	Cd 100 49.1 s	Cd 101 1.2 m	Cd 102 5.5 m	Cd 103 7.3 m	Cd 104 57.7 m	Cd 105 55.5 m	Cd 106 1.25	Cd 107 6.5 h	Cd 108 0.83	Cd 109 462.6 d	Cd 110 12.49	Cd 111 49 m	Cd 112 24.13	Cd 113 12.22
Ag 96 5.1 s	Ag 97 19 s	Ag 98 46.7 s	Ag 99 195 s	Ag 100 2.3 m	Ag 101 11.3 m	Ag 102 3.1 s	Ag 103 8.8 m	Ag 104 23.3 m	Ag 105 72 m	Ag 106 41.29 s	Ag 107 44.3 s	Ag 108 119 s	Ag 109 316.9 d	Ag 110 26.5 s	Ag 111 63 s	Ag 112 3.12 h
Pd 95 14 s	Pd 96 2.0 m	Pd 97 3.1 m	Pd 98 17.7 m	Pd 99 21.4 m	Pd 100 3.7 d	Pd 101 8.47 h	Pd 102 1.02	Pd 103 16.96 d	Pd 104 11.14	Pd 105 22.33	Pd 106 27.33	Pd 107 31.5 s	Pd 108 26.46	Pd 109 1.63 m	Pd 110 13.43	Pd 111 11.72
Rh 94 78.5 s	Rh 95 1.58 m	Rh 96 3.9 m	Rh 97 44 m	Rh 98 33 m	Rh 99 4.7 h	Rh 100 47 m	Rh 101 4.8 s	Rh 102 2.3 s	Rh 103 56.1 m	Rh 104 4.8 m	Rh 105 45 s	Rh 106 3.2 s	Rh 107 21.7 m	Rh 108 5.9 m	Rh 109 80 s	Rh 110 27.1 s
Ru 93 108 s	Ru 94 51.8 m	Ru 95 1.65 h	Ru 96 5.52	Ru 97 2.9 d	Ru 98 1.88	Ru 99 12.7	Ru 100 12.6	Ru 101 17.0	Ru 102 31.6	Ru 103 38.35 d	Ru 104 18.7	Ru 105 4.44 h	Ru 106 373.6 d	Ru 107 3.07 m	Ru 108 4.5 m	Ru 109 34.5 s
Tc 92 4.4 m	Tc 93 43.5 s	Tc 94 53 m	Tc 95 50 s	Tc 96 82.2 s	Tc 97 4.2 · 10 ⁶ a	Tc 98 6.0 m	Tc 99 15.8 s	Tc 100 14.2 m	Tc 101 13.3 m	Tc 102 54.2 s	Tc 103 21.2 s	Tc 104 18.2 m	Tc 105 7.6 m	Tc 106 36 s	Tc 107 21.2 s	Tc 108 5.17 s
Mo 91 85 s	Mo 92 14.84	Mo 93 4.87 s	Mo 94 9.25	Mo 95 15.92	Mo 96 16.68	Mo 97 9.55	Mo 98 24.13	Mo 99 66.0 h	Mo 100 1.15 · 10 ⁶ a	Mo 101 14.6 m	Mo 102 11.2 m	Mo 103 67.5 s	Mo 104 1.0 m	Mo 105 35.6 s	Mo 106 8.7 s	Mo 107 3.5 s
Nb 90 152 s	Nb 91 16.13 s	Nb 92 10.19 s	Nb 93 6.25 m	Nb 94 2.10 ⁶ a	Nb 95 86.6 h	Nb 96 23.4 h	Nb 97 53 s	Nb 98 31.8 s	Nb 99 2.5 m	Nb 100 3.1 s	Nb 101 7.1 s	Nb 102 43 s	Nb 103 1.5 s	Nb 104 0.8 s	Nb 105 2.95 s	Nb 106 1.0 s
Zr 89 4.19 m	Zr 90 51.45	Zr 91 11.22	Zr 92 17.15	Zr 93 1.5 · 10 ⁶ a	Zr 94 17.38	Zr 95 64.0 d	Zr 96 2.80	Zr 97 16.8 h	Zr 98 30.7 s	Zr 99 2.1 s	Zr 100 2.1 s	Zr 101 2.1 s	Zr 102 2.9 s	Zr 103 1.3 s	Zr 104 1.2 s	Zr 105 ~ 1 s
Y 88 105.6 d	Y 89 16.8 m	Y 90 3.19 h	Y 91 49.7 m	Y 92 3.54 h	Y 93 10.1 h	Y 94 18.7 m	Y 95 10.3 m	Y 96 9.6 s	Y 97 3.2 s	Y 98 2.9 s	Y 99 1.47 s	Y 100 9.4 s	Y 101 448 ms	Y 102 3.97 s	Y 103 1.79 s	Y 104 1.17 s
50	4,764	5,835	5,866	5,979	6,300	6,469	6,545	6,270	5,971	5,753	6,161	6,199	5,116	4,271	3,016	



Capture studies: Fe, Ni, Zn, and Se

Motivations:

- Study of the weak s-process component (nucleosynthesis up to $A \sim 90$)
- Contribution of massive stars (core He-burning phase) to the s-process nucleosynthesis.
- s-process efficiency due to bottleneck cross sections (Example: ^{62}Ni)





In addition:

Fe and Ni are the most important structural materials for nuclear technologies. Results of previous measurements at n_TOF show that capture rates for light and intermediate-mass isotopes need to be revised.

Capture studies: Fe, Ni, Zn, and Se


- Setup: C_6D_6 in EAR-1
- All samples are stable(*) and non-hazardous
- Metal samples preferable (oxides acceptable)



(*) except ^{79}Se

Capture studies: $A \approx 150$

- EAR-2 required
- Sample from ISOLDE?



- branching isotope in the Sm-Eu-Gd region: test for low-mass TP-AGB
- branching ratio (capture/ β -decay) provides infos on the thermodynamical conditions of the s-processing (if accurate capture rates are known!)

Capture studies: actinides


Neutron cross section measurements for nuclear waste transmutation and advanced nuclear technologies


$^{241,243}\text{Am}$	The most important neutron poison in the fuels proposed for transmutation scenarios. Build up of Cm isotopes.
$^{239,240,242}\text{Pu}$	(n, γ) and (n,f) with active canning. Build up of Am and Cm isotopes.
^{245}Cm	No data available.
$^{235,238}\text{U}$	Improvement of standard cross sections.
$^{232}\text{Th}, ^{233,234}\text{U}$ $^{231,233}\text{Pa}$	Th/U advanced nuclear fuels. ^{233}U fission with active canning.

All measurements can be done in EAR-1 (except ^{241}Am and ^{233}Pa)

Capture studies: actual TAC setup

Actual setup with thick sample cannings






Neutron
Absorber


^{10}B loaded
Carbon Fibre
Capsules

Neutron Beam


Capture studies: actual TAC setup



Capture studies: Low neutron sensitivity setup



Capture studies: active canning for simultaneous (n,γ) & (n,f) measurements





[<< back](#)

Measurement of capture cross sections of fissile materials (veto) and measurement of the $(n,\gamma)/(n,f)$ ratio.

Fission studies

Fission studies

absolute $^{235}\text{U}(n,f)$ cross section from (n,p) scattering




(n,p) larger or comparable up to 100 MeV

Beam	capture mode (2 mm \varnothing)
Scattering angle	30°
Target thickness	250 $\mu\text{g}/\text{cm}^2$
Detector radius	20 mm
Target-to-detector distance	250 mm

Fission studies

FF distributions in vibrational resonances




Principles:

- Time-tag detector for the “start” signal
- Masses (kinetic energies) of FF from position-sensitive detectors (MICROMEGAS or semiconductors)

Fission studies

cross sections with PPAC detectors: present setup



Measurements:



- $^{231}\text{Pa}(n,f)$
- Fission fragments angular distributions (45° tilted targets) for ^{232}Th , ^{238}U and other low-activity actinides

EAR-2 boost:

- measurements of $^{241,243}\text{Am}$ (in class-A lab)
- measurements of ^{241}Pu and ^{244}Cm (in class-A lab)

[<< back](#)

Fission studies with twin ionization chamber



Twin ionization detector with measurement of both FF (PPAC principle)


Measurements:

- FF yields: mass & charge
- Test measurement with ^{235}U then measurements of other MA

[<< back](#)

(n,p) , (n,α) & (n,lcp) measurements

1. CIC: compensated ion chamber already tested at n_TOF



For n_TOF-Ph2:

- four chambers in the same volume for multi-sample measurements

Measurements:

- $^{147}\text{Sm}(n,\alpha)$ (tune up experiment)
- ^6LiF target for calibration


EAR-2 boost:

- approx 100 times the ORELA count rate expected
- ^{67}Zn and ^{99}Ru (n,α) measurements

(n,p) , (n,α) & (n,lcp) measurements

2. MICROMEAS

already used for measurements of nuclear recoils at n_TOF




For n_TOF-Ph2:

- converter replaced by sample
- expected count rate: 1 reaction/pulse ($\sigma=200$ mb, $\text{Ø}=5\text{cm}$, $1\mu\text{m}$ thick)

[<< back](#)

(n,p) , (n,α) & (n,lcp) measurements

3. Scattering chambers with ΔE -E or ΔE - ΔE -E telescopes



Setup: in parallel with fission detectors

- ✓ production cross sections $\sigma(E_n)$ for (n,xc)
- ✓ $c = p, \alpha, d$
- ✓ differential cross sections $d\sigma/d\Omega$, $d\sigma/dE$

Measurements:


- ^{56}Fe and ^{208}Pb (tune up experiment)
- Al, V, Cr, Zr, Th, and ^{238}U
- a few $\times 10^{18}$ protons/sample in fission mode

[<< back](#)

Neutron scattering reactions

Direct n + n scattering experiment not feasible!

Alternatively, interaction of two neutrons in the final state of a nuclear reaction. Examples of such reactions are:




Kinematic locus of the $n + {}^2\text{H} \rightarrow n + p + n$ reaction for:

$$E_n = 50 \text{ MeV}$$

$$\Theta_n = 20^\circ, \Phi_n = 0^\circ$$

$$\Theta_p = 50^\circ, \Phi_p = 180^\circ$$




Neutron incident energy 30 – 75 MeV
in 2.5 MeV bins

[<< back](#)


<< back

$^{232}\text{Th}(n,\gamma)$: n_TOF & GELINA




Source: L Leal, IAEA CRP meeting, December 2004

$^{237}\text{Np}(n,\gamma)$ at LANSCE




(Analysis by
E.I. Esch and
R. Reifarth)

$^{237}\text{Np}(n,\gamma)$ at LANSCE



(Analysis by
E.I. Esch and
R. Reifarth)




$^{237}\text{Np}(n,\gamma)$ at n_TOF



[< back](#)

Parallel Plate Avalanche Counters (PPACs)

- 20x20 cm²
- Isobutane gas 7 mbar
- HV 500-600 V
- 3 mm between electrodes
- 1 anode (a few ns signal width)
- Electrode thickness: 1.5 μm (Mylar+Al)
- Deposit thickness : 100-300 μg/cm²
- Backing thickness : 0.1 μm (Al)
- : 1.5 μm (Mylar)
- Fission event identification: T2 in coincidence with T1



IN2P3 (IPN Orsay)

- position-sensitive!